

AN ELEMENTARY PROOF OF THE CROSS THEOREM IN THE REINHARDT CASE

MAREK JARNICKI AND PETER PFLUG

ABSTRACT. We present an elementary proof of the cross theorem in the case of Reinhardt domains. The results illustrates the well-known interrelations between the holomorphic geometry of a Reinhardt domain and the convex geometry of its logarithmic image.

1. INTRODUCTION. MAIN RESULT.

The problem of continuation of separately holomorphic functions defined on a cross has been investigated in several papers, e.g. [Ber 1912], [Sic 1969a], [Sic 1969b], [Akh-Ron 1973], [Zah 1976], [Sic 1981], [Shi 1989], [Ngu-Sic 1991], [Ngu-Zer 1991], [Ngu-Zer 1995], [Ngu 1997], [Ale-Zer 2001], [Zer 2002] and may be formulated in the form of the following *cross theorem*.

Theorem 1.1. *Let $D_j \subset \mathbb{C}^{n_j}$ be a domain of holomorphy and let $A_j \subset D_j$ be a locally pluriregular set, $j = 1, \dots, N$, $N \geq 2$. Define the cross*

$$\mathbf{X} := \bigcup_{j=1}^N A_1 \times \cdots \times A_{j-1} \times D_j \times A_{j+1} \times \cdots \times A_N.$$

Let $f : \mathbf{X} \rightarrow \mathbb{C}$ be separately holomorphic, i.e. for any $(a_1, \dots, a_N) \in A_1 \times \cdots \times A_N$ and $j \in \{1, \dots, N\}$ the function

$$D_j \ni z_j \mapsto f(a_1, \dots, a_{j-1}, z_j, a_{j+1}, \dots, a_N) \in \mathbb{C}$$

is holomorphic. Then f extends holomorphically to a uniquely determined function $\hat{f}$ on the domain of holomorphy

$$\widehat{\mathbf{X}} := \left\{ (z_1, \dots, z_N) \in D_1 \times \cdots \times D_N : \sum_{j=1}^N h_{A_j, D_j}^*(z_j) < 1 \right\}, \quad (*)$$

where h_{A_j, D_j}^ is the upper regularization of the relative extremal function h_{A_j, D_j} , $j = 1, \dots, N$.*

Recall that $h_{A, D} := \sup\{u \in \mathcal{PSH}(D) : u \leq 1, u|_A \leq 0\}$.

Observe that in the case where A_j is open, $j = 1, \dots, N$, the cross $\mathbf{X}$ is a domain in $\mathbb{C}^n$ with $n := n_1 + \cdots + n_N$. Moreover, by the classical Hartogs lemma, every separately holomorphic function on $\mathbf{X}$ is simply holomorphic. Consequently, the formula (*) is nothing else as a description of the envelope of holomorphy of $\mathbf{X}$. Thus, it is natural to conjecture that in this case the formula (*) may be obtained

2000 *Mathematics Subject Classification.* 32D15, 32A07.

Key words and phrases. separately holomorphic function, cross theorem, Reinhardt domain. The research was partially supported by the DFG-grant 436POL113/103/0-2.

without the cross theorem machinery. Unfortunately, we do not know any such a simplification.

The aim of this note is to present an elementary geometric proof of Theorem 1.1 in the case where D_j is a Reinhardt domain and A_j is a non-empty Reinhardt open set, $j = 1, \dots, N$. The proof (§ 4) will be based on well-known interrelations between the holomorphic geometry of a Reinhardt domain and the convex geometry of its logarithmic image. Moreover, the cross theorem for the Reinhardt case may be taught in any lecture on Several Complex Variables; its proof needs only some basic facts for Reinhardt domains (see [Jar-Pfl 2008]).

2. CONVEX GEOMETRY.

We begin with some elementary results related to the convex domains in $\mathbb{R}^n$.

Definition 2.1. Let $\emptyset \neq S \subset U \subset \mathbb{R}^n$, where U is a convex domain. Define the *convex extremal function*

$$\Phi_{S,U} := \sup\{\varphi \in \mathcal{CVX}(U), \varphi \leq 1, \varphi|_S \leq 0\},$$

where $\mathcal{CVX}(U)$ stands for the family of all convex functions $\varphi : U \rightarrow [-\infty, +\infty)$.

- Remark 2.2.** (a) $\Phi_{S,U} \in \mathcal{CVX}(U)$, $0 \leq \Phi_{S,U} < 1$, and $\Phi_{S,U} = 0$ on S .
 (b) $\Phi_{\text{conv}(S),U} \equiv \Phi_{S,U}$.
 (c) If $\emptyset \neq S_k \subset U_k \subset \mathbb{R}^n$, U_k is a convex domain, $k \in \mathbb{N}$, $S_k \nearrow S$, and $U_k \nearrow U$, then $\Phi_{S_k,U_k} \searrow \Phi_{S,U}$.
 (d) For $0 < \mu < 1$, let $U_\mu := \{x \in U : \Phi_{S,U}(x) < \mu\}$ (observe that U_μ is a convex domain with $S \subset U_\mu$). Then $\Phi_{S,U_\mu} = (1/\mu)\Phi_{S,U}$ on U_μ .

Indeed, the inequality “ $\geq$ ” is obvious. To prove the opposite inequality, let

$$\varphi := \begin{cases} \max\{\Phi_{S,U}, \mu\Phi_{S,U_\mu}\} & \text{on } U_\mu \\ \Phi_{S,U} & \text{on } U \setminus U_\mu \end{cases}.$$

Then $\varphi \in \mathcal{CVX}(U)$, $\varphi < 1$, and $\varphi = 0$ on S . Thus $\varphi \leq \Phi_{S,U}$ and hence $\Phi_{S,U_\mu} \leq (1/\mu)\Phi_{S,U}$ in U_μ .

- (e) Let $\emptyset \neq S_j \subset U_j \subset \mathbb{R}^{n_j}$, where U_j is a convex domain, $j = 1, \dots, N$, $N \geq 2$. Put

$$W := \left\{ (x_1, \dots, x_N) \in U_1 \times \dots \times U_N : \sum_{j=1}^N \Phi_{S_j,U_j}(x_j) < 1 \right\}$$

(observe that W is a convex domain with $S_1 \times \dots \times S_N \subset W$). Then

$$\Phi_{S_1 \times \dots \times S_N, W}(x) = \sum_{j=1}^N \Phi_{S_j,U_j}(x_j), \quad x = (x_1, \dots, x_N) \in W.$$

Indeed, the inequality “ $\geq$ ” is obvious. To prove the opposite inequality we use induction on $N \geq 2$.

Let $N = 2$. To simplify notation write $A := S_1$, $U := U_1$, $B := S_2$, $V := U_2$. Observe that $T := (A \times V) \cup (U \times B) \subset W$ and directly from the definition we get

$$\Phi_{A \times B, W}(x, y) \leq \Phi_{A,U}(x) + \Phi_{B,V}(y), \quad (x, y) \in T.$$

Fix a point $(x_0, y_0) \in W \setminus T$. Let

$$\mu := 1 - \Phi_{A,U}(x_0) \in (0, 1], \quad V_\mu := \{y \in V : \Phi_{B,V}(y) < \mu\},$$

$$\varphi := \frac{1}{\mu}(\Phi_{A \times B, W}(x_0, \cdot) - \Phi_{A,U}(x_0)).$$

Then φ is a well-defined convex function on V_μ , $\varphi < 1$ on V_μ , and $\varphi \leq 0$ on B . Thus, by (d), $\varphi(y_0) \leq \Phi_{B,V_\mu}(y_0) = \frac{1}{\mu}\Phi_{B,V}(y_0)$, which finishes the proof.

Now, assume that the formula is true for $N-1 \geq 2$. Put $S' := S_1 \times \cdots \times S_{N-1}$,

$$W' := \{(x_1, \dots, x_{N-1}) \in U_1 \times \cdots \times U_{N-1} : \sum_{j=1}^{N-1} \Phi_{S_j, U_j}(x_j) < 1\}.$$

Then, by the inductive hypothesis, we have

$$\Phi_{S', W'}(x') = \sum_{j=1}^{N-1} \Phi_{S_j, U_j}(x_j), \quad x' = (x_1, \dots, x_{N-1}) \in W'.$$

Consequently,

$$W = \{(x', x_N) \in W' \times U_N : \Phi_{S', W'}(x') + \Phi_{S_N, U_N}(x_N) < 1\}.$$

Hence, using the case $N = 2$ (to $S' \subset W'$ and $S_N \subset U_N$), we get

$$\Phi_{S_1 \times \cdots \times S_N, W}(x) = \Phi_{S', W'}(x') + \Phi_{S_N, U_N}(x_N) = \sum_{j=1}^N \Phi_{S_j, U_j}(x_j),$$

$$x = (x', x_N) = (x_1, \dots, x_N) \in W.$$

Notice that properties (d) and (e) correspond to analogous properties of the relative extremal function — cf. e.g. [Sic 1981].

Proposition 2.3. *Let $\emptyset \neq S_j \subset U_j \subset \mathbb{R}^{n_j}$, where U_j is a convex domain and $\text{int } S_j \neq \emptyset$, $j = 1, \dots, N$, $N \geq 2$, and define the cross*

$$T := \bigcup_{j=1}^N S_1 \times \cdots \times S_{j-1} \times U_j \times S_{j+1} \times \cdots \times S_N.$$

Then

$$\text{conv}(T) = \left\{ (x_1, \dots, x_N) \in U_1 \times \cdots \times U_N : \sum_{j=1}^N \Phi_{S_j, U_j}(x_j) < 1 \right\} =: W. \quad (1)$$

Proof. We may assume that S_j is convex, $j = 1, \dots, N$ (cf. Remark 2.2(b)). The inclusion “ $\subset$ ” is obvious. Let

$$T_j := S_1 \times \cdots \times S_{j-1} \times U_j \times S_{j+1} \times \cdots \times S_N, \quad j = 1, \dots, N,$$

$$T' := \bigcup_{j=1}^{N-1} S_1 \times \cdots \times S_{j-1} \times U_j \times S_{j+1} \times \cdots \times S_{N-1}, \quad S' := S_1 \times \cdots \times S_{N-1}.$$

(¹) It seems to us that this “convex cross theorem” is so far nowhere in the literature.

Recall (cf. [Roc 1972], Theorem 3.3) that

$$\begin{aligned} \text{conv}(T) \\ = \bigcup_{\substack{t_1, \dots, t_N \geq 0 \\ t_1 + \dots + t_N = 1}} t_1 T_1 + \dots + t_N T_N = \text{conv}((\text{conv}(T') \times S_N) \cup (S' \times U_N)). \quad (**) \end{aligned}$$

We use induction on N .

$N = 2$: To simplify notation write $A := S_1$, $U := U_1$, $p := n_1$, $B := S_2$, $V := U_2$, $q := n_2$. Using Remark 2.2(c), we may assume that U, V are bounded.

Since $\text{conv}(T)$ is open and $\text{conv}(T) \subset W$, we only need to show that for every $(x_0, y_0) \in \partial(\text{conv}(T)) \cap (U \times V)$ we have $\Phi_{A,U}(x_0) + \Phi_{B,V}(y_0) = 1$. Since U, V are bounded, we have $\overline{\text{conv}(T)} = \text{conv}(\overline{T})$ (cf. [Roc 1972], Theorem 17.2) and therefore, $(x_0, y_0) = t(x_1, y_1) + (1-t)(x_2, y_2)$, where $t \in [0, 1]$, $(x_1, y_1) \in \overline{A} \times \overline{U}$, $(x_2, y_2) \in \overline{U} \times \overline{B}$. First observe that $t \in (0, 1)$.

Indeed, suppose for instance that $(x_0, y_0) \in U \times (\overline{B} \cap V)$. Take an arbitrary $x_* \in \text{int } A$ and let $r > 0$, $\varepsilon > 0$ be such that the Euclidean ball $\mathbb{B}((x_*, y_0), r)$ is contained in $A \times V$ and $x_{**} := x_* + \varepsilon(x_0 - x_*) \in U$. Then

$$\begin{aligned} (x_0, y_0) &\in \text{int}(\text{conv}(\mathbb{B}((x_*, y_0), r) \cup \{(x_{**}, y_0)\})) \\ &\subset \text{int}(\text{conv}(\overline{T})) = \text{int}(\overline{\text{conv}(T)}) = \text{conv}(T); \end{aligned}$$

a contradiction.

Let $L : \mathbb{R}^p \times \mathbb{R}^q \longrightarrow \mathbb{R}$ be a linear form such that $L(x_0, y_0) = 1$ and $L \leq 1$ on T . Since $1 = L(x_0, y_0) = tL(x_1, y_1) + (1-t)L(x_2, y_2)$, we conclude that $L(x_1, y_1) = L(x_2, y_2) = 1$. Write $L(x, y) = P(x) + Q(y)$, where $P : \mathbb{R}^p \longrightarrow \mathbb{R}$, $Q : \mathbb{R}^q \longrightarrow \mathbb{R}$ are linear forms.

Put $P_C := \sup_C P$, $C \subset \mathbb{R}^p$, $Q_D := \sup_D Q$, $D \subset \mathbb{R}^q$. Since $L \leq 1$ on T and $L(x_1, y_1) = L(x_2, y_2) = 1$, we conclude that

$$\begin{aligned} P_A + Q_V &= 1, \\ P_U + Q_B &= 1. \end{aligned}$$

In particular, $P_A = P_U$ iff $Q_B = Q_V$. Consider the following two cases:

- $P_A < P_U$ and $Q_B < Q_V$: Then

$$\frac{P - P_A}{P_U - P_A} \leq \Phi_{A,U}, \quad \frac{Q - Q_B}{Q_V - Q_B} \leq \Phi_{B,V}.$$

Hence

$$\Phi_{A,U}(x_0) + \Phi_{B,V}(y_0) \geq \frac{P(x_0) - P_A}{1 - Q_B - P_A} + \frac{Q(y_0) - Q_B}{1 - P_A - Q_B} = 1.$$

- $P_A = P_U$ and $Q_B = Q_V$: Then $P_U + Q_V = 1$, which implies that $(x_0, y_0) \in U \times V \subset \{L < 1\}$; a contradiction.

Now, assume that the result is true for $N - 1 \geq 2$. In particular,

$$\text{conv}(T') = \left\{ (x_1, \dots, x_{N-1}) \in U_1 \times \dots \times U_{N-1} : \sum_{j=1}^{N-1} \Phi_{S_j, U_j}(x_j) < 1 \right\} =: W'.$$

Using (**), the case $N = 2$, and Remark 2.2(e), we get

$$\begin{aligned} \text{conv}(T) &= \text{conv}((W' \times S_N) \cup ((S' \times U_N))) \\ &= \{(x', x_N) \in W' \times U_N : \Phi_{S', W'}(x') + \Phi_{S_N, U_N}(x_N) < 1\} = W. \quad \square \end{aligned}$$

3. REINHARDT GEOMETRY.

Now we recall basic facts related to Reinhardt domains.

Definition 3.1. We say that a set $A \subset \mathbb{C}^n$ is a *Reinhardt set* if for every $(a_1, \dots, a_n) \in A$ we have

$$\{(z_1, \dots, z_n) \in \mathbb{C}^n : |z_j| = |a_j|, j = 1, \dots, n\} \subset A;$$

cf. [Jar-Pfl 2008], Definition 1.5.2. Put

$$\begin{aligned} \mathbf{V}_j &:= \mathbb{C}^{n-j-1} \times \{0\} \times \mathbb{C}^{n-j}, \quad \mathbf{V}_0 := \mathbf{V}_1 \cup \dots \cup \mathbf{V}_n, \\ \log A &:= \{(\log |z_1|, \dots, \log |z_n|) : (z_1, \dots, z_n) \in A \setminus \mathbf{V}_0\}, \quad A \subset \mathbb{C}^n, \\ \exp S &:= \{(z_1, \dots, z_n) \in \mathbb{C}^n \setminus \mathbf{V}_0 : (\log |z_1|, \dots, \log |z_n|) \in S\}, \quad S \subset \mathbb{R}^n, \\ A^* &:= \text{int}(\overline{\exp(\log A)}), \quad A \subset \mathbb{C}^n. \end{aligned}$$

We say that a set $A \subset \mathbb{C}^n$ is *logarithmically convex* (*log-convex*) if $\log A$ is convex; cf. [Jar-Pfl 2008], Definition 1.5.5.

Theorem 3.2 ([Jar-Pfl 2008], Theorem 1.11.13). *Let $\Omega \subset \mathbb{C}^n$ be a Reinhardt domain. Then the following conditions are equivalent:*

- (i) Ω is a domain of holomorphy;
- (ii) Ω is log-convex and $\Omega = \Omega^* \setminus \bigcup_{\substack{j \in \{1, \dots, n\} \\ \Omega \cap \mathbf{V}_j = \emptyset}} \mathbf{V}_j$.

Theorem 3.3 ([Jar-Pfl 2008], Theorem 1.12.4). *For every Reinhardt domain $\Omega \subset \mathbb{C}^n$ its envelope of holomorphy $\widehat{\Omega}$ is a Reinhardt domain.*

Corollary 3.4. *Let $\Omega \subset \mathbb{C}^n$ be a Reinhardt domain and let $\widehat{\Omega}$ be its envelope of holomorphy. Then*

- (a) $\mathbf{V}_j \cap \widehat{\Omega} = \emptyset$ iff $\mathbf{V}_j \cap \Omega = \emptyset$,
- (b) $\log \widehat{\Omega} = \text{conv}(\log \Omega)$.

Consequently, by Theorem 3.3,

$$\widehat{\Omega} = \text{int}(\overline{\exp(\text{conv}(\log \Omega))}) \setminus \bigcup_{\substack{j \in \{1, \dots, n\} \\ \Omega \cap \mathbf{V}_j = \emptyset}} \mathbf{V}_j =: \widetilde{\Omega}.$$

Proof. (a) If $\mathbf{V}_j \cap \Omega = \emptyset$, then the function $\Omega \ni z_j \mapsto 1/z_j$ is holomorphic on Ω . Thus, it must be holomorphically continuable to $\widehat{\Omega}$, which means that $\mathbf{V}_j \cap \widehat{\Omega} = \emptyset$.

(b) First observe that, by Remark 1.5.6(a) from [Jar-Pfl 2008], we get $\log \widetilde{\Omega} = \text{conv}(\log \Omega)$. Consequently, $\widetilde{\Omega}$ is a domain of holomorphy with $\Omega \subset \widetilde{\Omega}$. Hence, $\widehat{\Omega} \subset \widetilde{\Omega}$. Finally, $\log \Omega \subset \log \widehat{\Omega} \subset \log \widetilde{\Omega} = \text{conv}(\log \Omega)$. $\square$

Proposition 3.5 ([Jar-Pfl 2008], Proposition 1.14.20). *Let Ω be a log-convex Reinhardt domain.*

- (a) *Let $u \in \mathcal{PSH}(\Omega)$ be such that*

$$u(z_1, \dots, z_n) = u(|z_1|, \dots, |z_n|), \quad (z_1, \dots, z_n) \in \Omega.$$

Then the function

$$\log \Omega \ni (x_1, \dots, x_n) \mapsto u(e^{x_1}, \dots, e^{x_n})$$

is convex.

(b) Let $\varphi \in \mathcal{CVX}(\log \Omega)$. Then the function

$$\Omega \setminus \mathbf{V}_0 \ni z \mapsto \varphi(\log |z_1|, \dots, \log |z_n|)$$

is plurisubharmonic.

Corollary 3.6. Let $\emptyset \neq A \subset \Omega$, where Ω is log-convex Reinhardt domain and A is an Reinhardt open set. Then

$$h_{A,D}^*(z) = \Phi_{\log A, \log \Omega}(\log |z_1|, \dots, \log |z_n|), \quad z = (z_1, \dots, z_n) \in \Omega \setminus \mathbf{V}_0;$$

cf. Definition 2.1.

Proof. Since A and Ω are invariant under rotations, we easily conclude that

$$h_{A,D}^*(z) = h_{A,D}^*(|z_1|, \dots, |z_n|), \quad z = (z_1, \dots, z_n) \in \Omega.$$

Thus, by Proposition 3.5,

$$h_{A,D}^*(z) = \varphi(\log |z_1|, \dots, \log |z_n|), \quad z = (z_1, \dots, z_n) \in \Omega \setminus \mathbf{V}_0,$$

where $\varphi \in \mathcal{CVX}(\log \Omega)$. Clearly, $h_{A,D}^* = 0$ on A . Thus $\varphi = 0$ on $\log A$. Finally, $\varphi \leq \Phi_{\log A, \log \Omega}$.

To prove the opposite inequality, observe that by Proposition 3.5, the function

$$\Omega \setminus \mathbf{V}_0 \ni z \mapsto \Phi_{\log A, \log \Omega}(\log |z_1|, \dots, \log |z_n|)$$

is plurisubharmonic, $u < 1$, and $u = 0$ on $A \setminus \mathbf{V}_0$. Consequently, u extends to a $\tilde{u} \in \mathcal{PSH}(\Omega)$. Clearly, $\tilde{u} \leq 1$ and $\tilde{u} = 0$ on A . Thus $\tilde{u} \leq h_{A,D}^*$. $\square$

4. PROOF OF THE CROSS THEOREM (THEOREM 1.1) IN THE CASE WHERE D_j IS A REINHARDT DOMAIN OF HOLOMORPHY AND A_j IS AN OPEN REINHARDT SET, $j = 1, \dots, N$.

We have to prove that the envelope of holomorphy $\widehat{\mathbf{X}}$ of the domain $\mathbf{X}$ coincides with

$$\widetilde{\mathbf{X}} := \left\{ (z_1, \dots, z_N) \in D_1 \times \dots \times D_N : \sum_{j=1}^N h_{A_j, D_j}^*(z_j) < 1 \right\}.$$

First, observe that $\widetilde{\mathbf{X}}$ is a domain of holomorphy containing $\mathbf{X}$. Thus $\widehat{\mathbf{X}} \subset \widetilde{\mathbf{X}}$. On the other hand, by Proposition 2.3 and Corollary 3.6, $\log \widetilde{\mathbf{X}} = \text{conv}(\log \mathbf{X}) = \log \widehat{\mathbf{X}}$. Thus, using Corollary 3.4, we only need to show that if $\mathbf{V}_j \cap \widetilde{\mathbf{X}} \neq \emptyset$, then $\mathbf{V}_j \cap \mathbf{X} \neq \emptyset$. Indeed, let for example $a = (a_1, \dots, a_N) \in \mathbf{V}_n \cap \widetilde{\mathbf{X}} \neq \emptyset$. Take arbitrary $b_j \in A_j$, $j = 1, \dots, N-1$. Then $(b_1, \dots, b_{N-1}, a_N) \in \mathbf{V}_n \cap \mathbf{X}$. $\square$

REFERENCES

- [Akh-Ron 1973] N.I. Akhiezer, L.I. Ronkin, *On separately analytic functions of several variables and theorems on "the thin end of the wedge"*, Usp. Mat. Nauk. 28 (1973), 27–44.
- [Ale-Zer 2001] O. Alehyane, A. Zeriahi, *Une nouvelle version du théorème d'extension de Hartogs pour les applications séparément holomorphes entre espaces analytiques*, Ann. Polon. Math. 76 (2001), 245–278.
- [Ber 1912] S.N. Bernstein, *Sur l'ordre de la meilleure approximation des fonctions continues par des polynômes de degré donné*, Bruxelles, 1912.
- [Jar-Pfl 2008] M. Jarnicki, P. Pflug, *First Steps in Several Complex Variables: Reinhardt Domains*, European Mathematical Society Publishing House, 2008.
- [Ngu 1997] Nguyen Thanh Van, *Separate analyticity and related subjects*, Vietnam J. Math. 25 (1997), 81–90.

- [Ngu-Sic 1991] Nguyen Thanh Van, J. Siciak, *Fonctions plurisousharmoniques extrémales et systèmes doublement orthogonaux de fonctions analytiques*, Bull. Sci. Math. 115 (1991), 235–244.
- [Ngu-Zer 1991] Nguyen Thanh Van, A. Zeriahi, *Une extension du théorème de Hartogs sur les fonctions séparément analytiques* in Analyse Complexe Multivariables, Récents Développements, A. Meril (éd.), EditEl, Rende 1991, 183–194.
- [Ngu-Zer 1995] Nguyen Thanh Van, A. Zeriahi, *Systèmes doublement orthogonaux de fonctions holomorphes et applications*, Banach Center Publ. 31 (1995), 281–297.
- [Roc 1972] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, 1972.
- [Shi 1989] B. Shiffman, *On separate analyticity and Hartogs theorem*, Indiana Univ. Math. J. 38 (1989), 943–957.
- [Sic 1969a] J. Siciak, *Analyticity and separate analyticity of functions defined on lower dimensional subsets of $\mathbb{C}^n$* , Zeszyty Naukowe UJ 13 (1969), 53–70.
- [Sic 1969b] J. Siciak, *Separately analytic functions and envelopes of holomorphy of some lower dimensional subsets of $\mathbb{C}^n$* , Ann. Polon. Math. 22 (1969–1970), 147–171.
- [Sic 1981] J. Siciak, *Extremal plurisubharmonic functions in $\mathbb{C}^N$* , Ann. Polon. Math. 39 (1981), 175–211.
- [Zah 1976] V.P. Zahariuta, *Separately analytic functions, generalizations of Hartogs theorem, and envelopes of holomorphy*, Math. USSR-Sb. 30 (1976), 51–67.
- [Zer 2002] A. Zeriahi, *Comportement asymptotique des systèmes doublement orthogonaux de Bergman: une approche élémentaire*, Vietnam J. Math. 30 (2002), 177–188.

JAGIELLONIAN UNIVERSITY, INSTITUTE OF MATHEMATICS, ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND

E-mail address: Marek.Jarnicki@im.uj.edu.pl

CARL VON OSSIETZKY UNIVERSITÄT OLDENBURG, INSTITUT FÜR MATHEMATIK, POSTFACH 2503, D-26111 OLDENBURG, GERMANY

E-mail address: pflug@mathematik.uni-oldenburg.de